

<b>Rabbiteye Blueberry Varieties</b>				
<b>Cultivar</b>	<b>Chilling Requirements</b>	<b>Ripen Date (Southern MS)</b>	<b>Productivity</b>	<b>Vigor</b>
<b>Alapaha</b>	450-550 hours	late May to early June	VG	G
<b>Austin</b>	450-500 hours	early to late June	VG	M
<b>Brightwell</b>	350-400 hours	early June to early July	VG	G
<b>Climax</b>	400-450 hours	late May to early June	VG	G
<b>Columbus</b>	600 hours	late June to mid-July	VG	G
<b>DeSoto</b>	600-650 hours	mid-July to mid-Augus	VG	M
<b>Ira</b>	600 hours	mid-June to July	G	G
<b>Montgomery</b>	550 hours	late May to mid-June	VG	M
<b>Ocklocknee</b>	650-700 hours	early to late July	VG	G
<b>Onslow</b>	500-600 hours	late June to late July	VG	G
<b>Powderblue</b>	550-600 hours	late June to late July	G	G
<b>Premier</b>	500-550 hours	late May to early June	G	G
<b>Prince</b>	300-400 hours	mid-May to early June	VG	G
<b>Savory</b>	300-400 hours	late May to early June	G	G
<b>Tifblue</b>	600-700 hours	late June to July	G	G
<b>Vernon</b>	500-550 hours	early to late June	G	G
<b>Yadkin</b>	500-600 hours	late June to late July	G	G

**F=Fair, G=Good, VG=Very Good, P=Poor, L=Large, M=Medium, S=Small**